



Detection of *Cryptosporidium canis* in dog in Mosul city, Iraq

Eman D. Hadi, Israa A. Al- Robaiee

Department of Internal and Preventive Medicine, College of Veterinary Medicine, University of Mosul, Mosul, Iraq.

ARTICLE INFO.

Article history:

-Received: 1/ 11/ 2024

-Accepted: 24/ 12 / 2024

-Available online:

Keywords:

**Cryptosporidium canis,
Fluorescent**

Corresponding Author:

Name:

Israa A. Al- Robaiee

E-mail:

alrubaye_israa@yahoo.com

Tel:

ABSTRACT

The current study aimed to detect *Cryptosporidium canis* in dogs in Mosul, Iraq. To achieve this objective, fecal samples from a total of 95 local dogs located in Mosul city were collected between September 2023 and April 2024. Direct, flotation, and fluorescent techniques were used for identifying the parasite. Iodin and Giemsa stains used as basic protocols for fecal smear samples, alongside acridine orange stain was used for fluorescence microscopy monitoring. For *Cryptosporidium* species, 45 specimens (47.4%) had been confirmed. Dogs younger than one year old (59.6%), Male dogs (78.6%), stray dogs (56%) reported the most rates of Cryptosporidiosis. This study concluded that detection of *Cryptosporidium canis* with the fluorescent using (acridine orange) was the most effective technique that showed positive results.

1. Introduction

Cryptosporidiosis is a parasitic zoonotic disease linked to *Cryptosporidium* spp. This parasite is considered a gut protozoan, distributed around the world, and affects humans, pets, and dogs in both subclinical and clinical forms [1,2]. Canine can be infested with four spp. of *cryptosporidium* including: *C. canis*, *C. anderson*, *C. parvum* and *C.ubiquitum* [3], but the two prevalent species in dogs are *C. canis* and *C. parvum* [4]. Since 1981, over one hundred research projects studied *Cryptosporidium* infections in dogs from 27 countries globally. [2, 5].

When this protozoon infects the gastrointestinal tract, mostly in the ileal region, it can cause epithelial cell devastation, intestinal villi shrinking, poor absorption in the digestive system, with decreased gastric acid secretion and a deficiency in vitamin B12 [6,7] The majority of prominent clinical manifestations are anemia, Anorexia, loss of body weight and frequent diarrhea [7].

One of the two modes of how this protozoon could possibly transmitted: directly by fecal-oral rout or indirectly through ingestion contaminated food or water Furthermore, this protozoan may spore oocysts in the host's cell, promote self-infected among several hosts and be active when excretion via feces and survive in extreme conditions [7,8].

Plenty of GIT protozoa in dogs have been identified throughout Iraq in different places, such as Baghdad [9], Mosul [10,11], Sulaimani [12], Basra [13,14], Dohuk [15], and Erbil [16]. In Baghdad, the two prevalent species of infected dogs are *C. canis* and *C. parvum* [9]. In Mosul city, 42% of dogs and 52% of cats were infected with *Cryptosporidium* spp.[11]. As far as we are aware, there is limited information about the description of *Cryptosporidium canis* in dogs in Mosul. Therefore, detecting and identifying *Cryptosporidium canis* in dogs in Mosul, Iraq was the goal of this study.

2. Materials and Methods

2.1 Animal's sampling:

Fecal samples from 95 dogs (20 domesticated dogs and 75 stray dogs) were collected, including both sexes and diverse ages between September 2023 and April 2024 from the University Veterinary Hospital at the University of Mosul, the Veterinary Hospital in Al-Rashidiya, and other locations of Mosul city.

2.2 Laboratory examination:

Coprological Qualitative methods included the direct and the flotation using zinc sulphate solution. Fecal films were prepared and examined using a fluorescent microscope with acridine orange stain as well as Iodine and Giemsa stains, which was further investigated using an optical microscope at 10X and 40X resolution [17].

2.3 Statistical analysis:

Analysis of the data was conducted on SPSS (Version 17; SPSS Inc., Chicago, USA). $P < 0.05$ has been taken into account when calculating statistical importance.

3.Results and Discussion:

The existence of *Cryptosporidium* spp. was observed by microscopy of the oocyst's size and morphological appearance using direct method, flotation with zinc sulphate solution in addition to Giemsa and Iodine stains (Fig. 1). The identified oocysts have oval to spherical appearance containing four sporozoites with a blue backdrop (Fig. 1), but when observed under a fluorescent microscope, these protozoa appeared as greenish with acridine orange stain (Fig. 2)




Figure1: *Cryptosporidium* Spp. With Giemsa stain, Iodine using ocular micrometer under light Microscope 10X.




Figure 2: Fluorescing *Cryptosporidium* Spp. using acridine orange stain, examined under fluorescent microscope 20X.

A total of 45 dogs (47.4%) were positive for cryptosporidium oocysts, Puppies recorded a higher infection rate (52.9%) than older animals (Table.1). There were significant ($P<0.05$) variations in the percentage of

positive samples for the cryptosporidiosis according to the age (puppies 52.9%), the gender (males 78.6%), and stray dogs (56%, Table 1).

Table (1): Total infection rate of *Cryptosporidium* spp. in dogs according to age, gender and housing

Variable	No. of examined sample	No. of Positive	Percentage (%)
Age			
≤ 1 year (young)	34	18	52.9 ^a
>1 year (adult)	61	27	44.3 ^b
Gender			
Females	53	12	22.6 ^b
Males	42	33	78.6 ^a
Housing			
Stray	75	42	56 ^a
Domestic	20	3	15 ^b
Total	95	45	47.4

The different small letters refer to the presence of significantly different at ($P < 0.05$).

4. Discussion

This study demonstrated that *Cryptosporidium canis* total infection rate was 47.4% (45/95) in canine, and that's identical to a previous study published in Mosul city which concluded that 42% of dogs were infected with *Cryptosporidium* spp. [11], while this ratio was not in line with the data obtained in Baghdad [9]. In addition to documentation in Iran [18]. The outcomes opposed those of ours, with prevalence rates of 8%. The elevation in infection rate may be attributed to heavy contamination of water and food with oocyst as well as suitable environment for oocysts to persist for long periods, furthermore variation in methods of sampling and diagnostic techniques [19].

Major variations occurred across the researched dogs in the present research. The most frequently documented infection rate was 52.9% in young dogs in contrast to the ratio reported in adult animals. These outcomes line up with many research investigations published in Iraq by researchers [20] who documented that infection rate in puppies (69%) was higher than adults (31%).

Conclusion:

The present deduced that examine of *Cryptosporidium canis* with the fluorescent technique using (acridine orange) showed positive results. This technique is beneficial for examine *Cryptosporidium* spp. in various animals.

Conversely, in South Africa an elevated infection rate of *Cryptosporidium canis*. was seen in adults (19%) than in puppies (6%) [21], which disagrees with our study. There is a potential relationship between age and danger of infection with *Cryptosporidium* which may be attributed to the fact that young dogs have an incomplete immune system, in addition to malnutrition following weaning which makes them more susceptible to being infected [11].

The majority of male dogs appeared to have the highest percentage (78.6%) for cryptosporidiosis, while the minor percentage (22.6%) reported in the females which agrees with another research in Iraq [11,20]. In contrast, it disagreed with a documentation in Karbala province in Iraq which recorded the higher infection rate in female dogs 48.27% compared to male dogs 23.89% [22]. The present and previous research illustrates that both sex of dogs has an equal probability to be infected when exposed to contaminated substances [23]. Furthermore, the high prevalence of this protozoa documented in stray dogs was more common.

Acknowledgment:

The researchers thank for supporting regulated by veterinary teaching hospital at the College of Veterinary Medicine, University of Mosul, Mosul, Iraq.

Conflict of Interest:

It was approved by the authors and there was no conflict of interest.

References

- [1] Pumipuntu, N. and Piratae, S. (2018). Cryptosporidiosis: a zoonotic disease concern. *Vet. World*,11,681–6. doi:10.14202/ vet world.2018.681- 686.
- [2] Taghipour, A., Olfatifar, M., Bahadory, S., Godfrey, S.S., Abdoli, A. and Khatami, (2020). The global prevalence of *Cryptosporidium* infection in dogs: a systematic review and meta-analysis. *Vet Parasitol.*,28,109093. Doi: 10.1016/j.vetpar.2020.109093.
- [3] Rosanowski, S.M., Banica, M. and Ellis, E. (2018). The molecular characterization of *Cryptosporidium* species in relinquished dogs in Great Britain: a novel zoonotic risk? *Parasitol. Res.*, 117, 1663– 1667. doi: 10.1007/s00436-018-5857-z.
- [4] Shenquan, L., Xuhui, L., Yongxiang, S., Nanshan, Q., Minna, L., Caiyan, W., Juan, L., Junjing, H., Linzeng, Y., Haiming, C., Wenwan, X., Mingfei, S and Guoqing, L. (2020). Occurrence and genotypes of *Cryptosporidium* spp., *Giardia duodenalis*, and *Blastocystis* spp. in household, shelter, breeding, and pet market dogs in Guangzhou, southern China. *Scientific reports*. doi.org/V,10.1038/s41598-020-74299-P,1-11.
- [5] Tzipori, S. and Campbell, I. (1981) Prevalence of *Cryptosporidium* antibodies in 10 animal species. *J Clin Microbiol*,14,455–6. DOI: 10.1128/jcm.14.4.455-456.1981.
- [6] Nilson, B., Diego, A.G.L. and Regina, M.B.F. (2012): Vector-borne and zoonotic disease. *Parasitol*,12, 5-11.
- [7] Constable, P.D., Hinchcliff, K.W., Done, S.H. and Grunberg, W. (2017) *Veterinary Medicine. A textbook of the diseases of cattle, sheep, goats and horses* (11th ed), WB Saunders.
- [8] Kostopoulou, D., Claerebout, E., Arvanitis, D., Ligda, P., Voutzourakis, N. and Casaert, S. (2017). Abundance zoonotic potential and risk factors of intestinal parasitism amongst dog and cat populations: the scenario of Crete, Greece. *Parasit Vectors*,10,43-45. Doi: 10.1186/ s13071-017-1989-8
- [9] Faraj, A. (2019). Traditional and Molecular study of *Cryptosporidium* spp. in Domestic dogs in Baghdad city, Iraq. *The Iraqi Journal of Agricultural*,50(4),1094-1099.<https://doi.org/10.36103/ijas.v50i4.754>.
- [10] Al-Saeed, W.M., Al-Dabbagh, N.Y. and Mahmood, H.J. (2009) Serological study of toxocariasis in children in Mosul Province. *Med J Babylon*,6,455–464. <https://search.emarefa.net/detail/BIM-334021>.
- [11] Hadi, E. D., Suleiman, E. G., Al-Obadi, Q. T., and Arslan, S. H. (2014). Diagnostic study of *Cryptosporidium* spp. And *Giardia* spp. in stray dogs and cats in Mosul city, Iraq. *Iraqi Journal of Veterinary Sciences*, 28(1), 19-24. <https://doi.org/10.33899/ijvs.2014.89337>.
- [12] Mohammed, M.O. (2013) The human Seroprevalence of *Echinococcus Granulosus* in Sulaimani Governorate. *The Iraqi Postgraduate Medical Journal* 12:45–50. https://www.researchgate.net/publication/321807889_The_Human_Seroprevalence_of_Echinococcus_Granulosus_in_Sulaimani_Governorate.
- [13] Abu Tabeekh, M.A.S. and Thuwaini, M.M. (2015) Retrospective study of several zoonotic diseases affected human being in Basra governorate. *Mirror Res Vet Sci Anim* 4:8–16. <http://mirrorofresearchinveterinarysciencesandanimals.com/>.
- [14] Al-Jassim, K. B., Mahmmoud, Y. S., Salem, Z. M., and Al-Jubury, A. (2017). Epidemiological investigation of gastrointestinal parasites in dog populations in Basra province, Southern Iraq. *Journal of Parasitic Diseases*, 41(4), 1006-1013. DOI: 10.1007/s12639-017-0926-2.
- [15] Al-Saeed, A.T. and Issa, S.H. (2006). Frequency of *Giardia lamblia* among children in Dohuk, northern Iraq. *East Mediterr Health J*, 12,555–561. PMID: 17333793.
- [16] Saeed, I., Kapel, C., Saida, L.A., Willingham, L. and Nansen, P. (2000). Epidemiology of *Echinococcus granulosus* in Arbil province, northern Iraq, 1990–1998. *J*

Helminthol, 74,83–88.
10.1017/s0022149x00000111.

DOI: 181.<https://dx.doi.org/10.17582/journal.jahp/2024/12.s1.174.181>.

[17] Zajac, M., Conboy, G.A., Little, S.E. and Reichard, M.V. (2021). Veterinary Clinical Parasitology 9th ed. Wiley BlackWell LTd.

[18] Tavalla, M., Kord, E., Abdizadeh, R., Asgarian, F. (2017). Molecular study of Cryptosporidium spp. in dogs from southwest of Iran. Jundishapur J. Microbiol. 1-6. doi.org/10.5812/jjm.43412.

[19] Rasha, M.A. G., Abdallah, M.A. M, Ayman, A. S. and Amany, M. A.E. (2018). Molecular Screening and Genotyping of Cryptosporidium Species in Household Dogs and In-Contact Children in Egypt: Risk Factor Analysis and Zoonotic Importance. vector-borne and zoonotic diseases, 1-9, DOI: 10.1089/vbz.2017.2254.

[20] Al-Asady, S.I.J, Al-Hasnawy, M.H. (2024). Investigation of Cryptosporidium species infecting dogs. J. Anim. Health Prod. 12, 174-

[21] Amidou, S., Machuene, A.T. & Pascal, B. (2013). The epidemiology of Cryptosporidium in cats and dogs in the Thohoyandou region, South Africa. African Journal of Microbiology Research, 7(21), 2510-2518. DOI:10.5897/AJMR12.1391.

[22] Al-Yasary, J.T., Faraj, A.A. (2021). Genotyping of Cryptosporidium spp. in domestic dogs and humans in Karbala Province Iraq. Annls Roman. Soc. Cell Biol., 25(6),4120-4125.<http://annalsofrscb.ro/index.php/journal/article/view/6199>.

[23] Olabanji, G.M., Maikai, B.V., Otolurin, G.R. (2016). Prevalence and risk factors associated with faecal shedding of Cryptosporidium oocysts in dogs in the federal capital territory, Abuja, Nigeria. Vet. Med. Int., 2016,4591238. <https://doi.org/10.1155/2016/4591238>.

الكشف عن طفيلي الخبيئات البوغية الكلبية في الكلاب في الموصل، العراق

ايمان دحام هادي، اسراء عبد الغني

جامعة الموصل ، كلية الطب البيطري، فرع الطب الباطني والوقائي، الموصل، العراق

الملخص

هدفت الدراسة الحالية إلى الكشف عن طفيلي الخبيئات البوغية الكلبية في الكلاب في مدينة الموصل /العراق. ولتحقيق هذا الهدف، تم جمع عينات براز من إجمالي 95 كلبًا محليًا خلال الفترة من ايلول 2023 الى نيسان 2024. وتم استخدام الطريقة المباشرة وطريقة الطفو والتألق المناعي لتحديد الطفيلي فضلًا عن استخدام صبغة اليود وصبغة كيمزا كبروتوكولات أساسية لمسحات البراز، إلى جانب استخدام صبغة البرتقالي الأكريدين للفحص بالمجهر المتألق. إذ تم تأكيد 45 عينة على أنها موجبة لطفيل الخبيئات البوغية الكلبية وبنسبة إصابة كلبية (47.4%). وسجلت أعلى معدلات الإصابة في كل من الكلاب التي يقل عمرها عن عام واحد (59.6%)، وجنس الكلاب من الذكور (78.6%) والكلاب السائبة (56%). نستنتج من هذه الدراسة بأن الكشف عن طفيلي *Cryptosporidium canis* بتقنية التألق المناعي باستخدام صبغة الاكردين البرتقالية كان أكثر التقنيات فعالية التي أظهرت نتائج إيجابية.

الكلمات الدلالية: الخبيئات البوغية الكلبية، التألق المناعي